

Bravelle (urofollitropin) side effects

Get emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat. Some women using this medicine have developed a condition called ovarian hyperstimulation syndrome (OHSS). This syndrome may be more likely to occur within the first 10 days after receiving the last dose in a treatment cycle. OHSS can be a life-threatening condition. Stop using urofollitropin and call your doctor at once if you have any symptoms of OHSS:

- severe pelvic pain;
- swelling of your hands or legs;
- stomach pain and swelling;
- shortness of breath;
- weight gain;
- nausea, vomiting, diarrhea; or
- urinating less than usual.

Less serious side effects may include:

- mild stomach upset, constipation;
- mild pelvic pain, cramps;
- breast tenderness;
- skin rash;
- hot flashes;
- acne; or
- pain, swelling, redness, itching, or irritation where the medicine was injected.

This is not a complete list of side effects and others may occur. Tell your doctor about any unusual or bothersome side effect.

What other drugs will affect Bravelle (urofollitropin)?

There may be other drugs that can interact with urofollitropin. Tell your doctor about all your prescription and over-the-counter medications, vitamins, minerals, herbal products, and drugs prescribed by other doctors. Do not start a new medication without telling your doctor.